

Points in the SLHC Collection:


Match the type of point to the timeline

- Kirk/Palmer Notched Point
- Fox Creek Points
- Poplar Island Points
- Bifurcate Points
- Pequea Points
- Basal-Notched Points
- Amos Point
- Teardrop Points
- Various Broadspear Types
- Fishtail Points
- Dalton-type Point
- Fluted point
- Triangle Points
- Jacks Reef Points
- Vosburg Points




- Approximately 13,000 years ago
- Between 9500 and 9000 B.C.
- Approx. 9000 B.C. to 8500 B.C.
- Approx. 6500 B.C. to 6000 B.C.
- Approx. 6500 B.C. to 3000 B.C.
- Approx. 5500 B.C. to 200 B.C.
- Approximately 2500 B.C. to 1500 B.C.
- 1000 B.C. to 0
- 0 to 1000 A.D.
- Approx. 0 to 1000 A.D.
- Approximately 0 to 1550 A.D.
- Approx. 1000 A.D. to 1550 A.D.
- 8000 B.C. to 6500 B.C.
- 8000 B.C. to 6500 B.C.
- 5000 B.C. to 1000 A.D.
- 1200 B.C. to 1000 A.D.


Illustrate the type of point



Fishtail Points




Basal-Notched Points



Triangle Points



Bifurcate Points



Dalton-type Point



Jacks Reef Points

points would have been used for: _____
 types of rock made into a point: _____